



**QUEEN'S
UNIVERSITY
BELFAST**

Abdominal Aortic Aneurysm Genetic Associations: Mostly False? A Systematic Review and Meta-analysis

Bradley, D. T., Badger, S. A., McFarland, M., & Hughes, A. E. (2016). Abdominal Aortic Aneurysm Genetic Associations: Mostly False? A Systematic Review and Meta-analysis. *European Journal of Vascular Endovascular Surgery*, 51(1), 64-75. <https://doi.org/10.1016/j.ejvs.2015.09.006>

Published in:
European Journal of Vascular Endovascular Surgery

Document Version:
Peer reviewed version

Queen's University Belfast - Research Portal:
[Link to publication record in Queen's University Belfast Research Portal](#)

Publisher rights

© 2016 Elsevier Ltd.

This manuscript version is made available under the CC-BY-NC-ND 4.0 license <http://creativecommons.org/licenses/by-nc-nd/4.0/> which permits distribution and reproduction for non-commercial purposes, provided the author and source are cited.

General rights

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact openaccess@qub.ac.uk.

Title: Abdominal Aortic Aneurysm Genetic Associations: Mostly False? A Systematic Review and Meta-Analysis

Declan T Bradley^a dbradley09@qub.ac.uk (corresponding)

Stephen A Badger^b stephenbadger@btinternet.com

Marie McFarland^c marie.mcfarland@belfasttrust.hscni.net

Anne E Hughes^{a1} annehughes169@btinternet.com

- a. Centre for Public Health, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Institute of Clinical Sciences, Block B, Royal Victoria Hospital, Belfast BT12 6BA, UK
 - b. Mater Misericordiae University Hospital, Eccles Street, Dublin, Ireland
 - c. Department of Pathology, Institute of Pathology Building, Royal Victoria Hospital, Belfast Health and Social Care Trust, Grosvenor Road, Belfast, BT12 6BL
1. Recently retired, formerly of QUB.

Category: **Review**

Short title (50 characters): **AAA genetics: Systematic Review and Meta-analysis**

Word count including references: 4,995

Abstract

Background

Many associations between abdominal aortic aneurysm (AAA) and genetic polymorphisms have been reported. It is unclear which are genuine and which may be caused by type-1 errors, biases and flexible study design.

Objectives

To identify associations supported by current evidence and to investigate the effect of study design on reporting associations.

Data Sources

MEDLINE, Embase and Web of Science.

Review Methods

Reports were dual-reviewed for relevance and inclusion against pre-defined criteria (studies of genetic polymorphisms and AAA risk). Study characteristics and data were extracted using an agreed tool and reports assessed for quality. Heterogeneity was assessed using I^2 and fixed- and random-effects meta-analyses were conducted for variants that were reported at least twice, if any had reported an association. Strength of evidence was assessed using a standard guideline.

Results

Searches identified 467 unique articles, of which 97 were included. Of 97 studies, 63 reported at least one association. Of 92 studies that conducted multiple tests, only 27% corrected their analyses. Two hundred and sixty three genes were investigated

and associations were reported in polymorphisms in 87 genes. Associations in *CDKN2BAS*, *SORT1*, *LRP1*, *IL6R*, *MMP3*, *AGTR1*, *ACE* and *APOA1* were supported by meta-analyses.

Conclusions

Uncorrected multiple testing and flexible study design (particularly testing many inheritance models and subgroups, and failure to check for Hardy-Weinberg Equilibrium) contributed to apparently false associations being reported. Heterogeneity, possibly due to the case-mix, geographical, temporal and environmental variation between different studies, was evident. Polymorphisms in nine genes had strong or moderate support on the basis of the literature at this time. We make suggestions for improving AAA genetics study design and conduct.

Key words

Aortic Aneurysm, Abdominal

Genetics

Polymorphism, Single Nucleotide

Review [Publication Type]

Meta-Analysis [Publication Type]

It has been claimed that most published research findings are false and that studies of complex genetic diseases are especially prone to reporting false associations.¹ Abdominal aortic aneurysms (AAA) occur in Mendelian disorders,² but are usually multifactorial. Understanding AAA pathogenesis might lead to new treatments or preventive measures. No meaningful personalised genetic risk prediction³ has been realised for AAA.

AAA is usually defined as infra-renal aortic diameter $\geq 30\text{mm}$.⁴ Using alternative definitions (infrarenal aortic diameter:suprarenal aortic diameter ratio, or comparison to predicted diameter) has striking effects on prevalence.⁵ Participants may be recruited from screening programmes or surgical departments, affecting case-mix. The prevalence of old age, tobacco use and medication use has changed over time and differs between locations, so gene-environment interactions could affect results.⁶⁻⁸

Ioannidis claimed that six factors increased the likelihood of research findings being false: small study size; small effect size; multiple testing; flexible study design; financial or other interests and the 'hotness' of the field.¹ The *STrengthening the REporting of Genetic Association Studies (STREGA)* guidance aims to make reports consistent and transparent,⁹ but there is variation in the analytic and statistical methods used. This study aimed to identify gene associations supported by evidence and to investigate the role of study design in reporting associations.

Methods

Search Strategy

We searched databases using MeSH terms (Table 1; Supplementary Methods) on 25 August 2014. The Medline search was simply “Aortic Aneurysm, Abdominal/genetics”, limited to human studies. The National Institutes of Health association database, references and relevant journals were also searched.

Inclusion Criteria

- Population: cases with AAA (infra-renal aorta >30mm, infra-renal/supra-renal aortic ratio ≥ 1.5 , surgical repair or ruptured AAA); controls were untested individuals, tested controls or hospital controls.
- Exposure: germline genomic polymorphisms.
- Comparison: disease risk in individuals with/without polymorphisms or odds of exposure in individuals with/without AAA in case-control, cross-sectional or cohort studies.
- Outcome: presence/absence of AAA.

Exclusion Criteria

Studies with single-disease controls (e.g. ischaemic heart disease), studies of somatic mutations, linkage studies and studies of AAA growth or size were excluded.

Study Selection and Data Abstraction

Two authors independently assessed titles and abstracts for relevance and full reports for inclusion. Two authors independently extracted quality and genotype/risk data for polymorphisms reported ≥ 2 times with an association reported at least once. Additional characteristics were abstracted by one author

Data analysis

Median study size was calculated for each year. Odds ratios for heterozygotes and homozygotes were calculated. Where genotype counts were not available, inheritance models from original reports were used. Additive model analyses were conducted using binary logistic regression in IBM SPSS Statistics v19.0 (IBM Corp, Armonk, NY), with genotypes coded as 0, 1 or 2. We re-analysed original *CRP* data from Badger *et al.*¹⁰ to allow meta-analysis with the results of Saratzis *et al.*¹¹ Hardy Weinberg Equilibrium (HWE) deviation was assessed by Haldane's Exact test in *HardyWeinberg* in *R*. Sensitivity analyses excluded studies with HWE deviation ($p < 0.05$). SNP Annotation and Proxy Search (Broad Institute, Cambridge, Massachusetts) identified markers with $r^2 > 0.9$. Where study data were published more than once (i.e. a study was updated or reported study populations overlapped), the selection of the report for inclusion was based on the reporting of genotype counts/frequencies or risk in preference to allelic or specific inheritance models. If the mode of reporting was the same then the larger study was included.

Heterogeneity index (I^2) and Cochrane's Q were estimated using *meta* in *R*. $I^2 > 50\%$ was large, and 25-50% moderate.¹² Fixed and random effects inverse variance-weighted meta-analysis was conducted using *meta* in *R*. Significance was $p < 0.05$ in fixed effects meta-analysis if $I^2 < 25\%$ or random effects meta-analysis if $I^2 \geq 25\%$.

Results

Search results

Searches yielded 614 records. Ten were identified from other sources. There were 477 unique records. Ninety seven were included (Figure 1). Excluded studies are listed (Supplementary Table 1).

Study types and reports of associations

There were 91 candidate gene studies, four genome-wide association studies (GWAS) and two genetic risk score (GRS) studies (Table 2). Fifty six candidate studies, both GRS and all GWAS reported associations.

Study size and frequency

Median study size for a candidate gene or GRS study was 710 participants (range 91-83,024). Study size and frequency of reports increased over time (Figure 2).

There was a median of three reports/year between 1994 and 25 August 2014.

Study quality

Quality indicators (Table 2) and detailed characteristics (Supplementary Table 2) are shown. Sixty two studies (64%) reported an association. Ninety two studies reported multiple tests and 25 (27%) adjusted analyses for this.

Genes investigated

Two hundred and sixty three genes were investigated (Supplementary Table 3), with 87 associated at least once. The median study size for reports that claimed an association was 811, and that of those that did not was 710 (Independent Samples Mann-Whitney U standardised test statistic=0.10, $p=0.92$).

Selection of polymorphisms for meta-analysis

We identified genes that featured in at least two studies and that had at least one association with sufficient data (Supplementary Table 3; Supplementary Table 4).

Results of meta-analyses

Meta-analyses are summarised (Table 3) and shown in full (Supplementary Table 5). Associations between AAA and 9p21 rs10757278, *SORT1* rs599839, *LRP1* rs1466535, *MMP3* rs3025058, *AGTR1* rs5186, *ACE* rs4646994 and *APOA1* rs964184 were supported. *CRP* rs3091244 lost significance in sensitivity analysis. *MMP2* rs243865 was associated but one of two studies deviated from HWE. Six were supported by strong evidence, three by moderate evidence and 25 had weak evidence (Table 4).

Discussion

Most studies reported associations, but only six were supported by strong evidence. Two were reported in genome-wide studies (*DAB2IP* and *LRP1*),^{72,78} and four in candidate gene studies (9p21/*CDKN2BAS*, *IL6R*, *LPA* and *SORT1*).^{50,93,94,108} There was moderate evidence for associations with *LDLR*, *MMP3* and *AGTR1* polymorphisms. *MMP3* rs3025058 was tentatively suggested to be an AAA risk factor in 1999 in a small study by Yoon *et al.* though it was not statistically significant after correction for multiple testing.¹⁸ Subsequent studies appear to support its effect in homozygosity. Other associations currently supported by moderate or strong evidence were all made since 2008.

Inter-study inconsistency may be due to heterogeneity of effect, systematic error or bias. Multiple testing was usually uncorrected and flexible analysis was common: Many studies tested numerous inheritance models and subgroups, reporting significance when $p < 0.05$. Several studies claimed HWE, but deviated significantly ($p < 0.05$).^{11,99}

We propose that it may be time for the genetic epidemiology research community to consider prospective registration of aetiological studies to avoid flexible *post hoc*

analyses. Simple standard steps, such as planning for adequate power, using contemporaneous controls and presenting counts and risk estimates for genotypes rather than assuming models of inheritance, checking frequencies in population databases, presenting analyses of HWE deviation and correcting for multiple testing would greatly improve the quality of reports in this field.

Our review excluded studies of aneurysm size or growth, and did not attempt to integrate results for 27 associated polymorphisms that had been reported only once.

Supported associations suggest the importance of lipoproteins. *LRP1*, *LDLR*, *SORT1* and the 9p21 locus affect cholesterol metabolism and atherosclerosis. *LRP1* has other important regulatory roles, including regulation of extracellular matrix breakdown by the endocytosis of proteinases.¹⁰⁹ *LPA* produces lipoprotein A which increases cardiovascular risk.¹¹⁰ *IL6R* polymorphisms alter cardiovascular risk, possibly through inflammation.¹¹¹⁻¹¹⁴ *MMP3* affects atherosclerosis and tissue remodelling.¹¹⁵ Its association in this review is in agreement with a recent meta-analysis.¹¹⁶ *AGTR1* affects blood pressure, which is consistent with the association between hypertension and AAA.^{6,117} *DAB2IP* is a tumour-suppressor gene involved in cell signalling, survival, migration, maturation and apoptosis.⁷² Understanding their roles may help develop prevention strategies based on understanding key biological pathways. Improving future study design will avoid wasteful false associations.

Acknowledgements

An earlier version of this work was presented at ESVES 2013. DTB was supported by a doctoral fellowship from the PHA.

References

1. Ioannidis JP. Why most published research findings are false. *PLoS medicine* 2005;2:e124.
2. Roberts WC, Honig HS. The spectrum of cardiovascular disease in the Marfan syndrome: a clinico-morphologic study of 18 necropsy patients and comparison to 151 previously reported necropsy patients. *Am Heart J* 1982;104:115-35.
3. Collins FS, Green ED, Guttmacher AE, Guyer MS, US National Human Genome Research, Institute. A vision for the future of genomics research. *Nature* 2003;422:835-47.
4. Steinberg CR, Archer M, Steinberg I. Measurement of the abdominal aorta after intravenous aortography in health and arteriosclerotic peripheral vascular disease. *Am J Roentgenol* 1965;95:703-8.
5. Wanhainen A. How to define an abdominal aortic aneurysm--influence on epidemiology and clinical practice. *Scand J Surg* 2008;97:105,9.
6. Kent KC, Zwolak RM, Egorova NN, Riles TS, Manganaro A, Moskowitz AJ, et al. Analysis of risk factors for abdominal aortic aneurysm in a cohort of more than 3 million individuals. *J Vasc Surg* 2010;52:539-48.
7. Singh K, Bonna KH, Jacobsen BK, Bjork L, Solberg S. Prevalence of and risk factors for abdominal aortic aneurysms in a population-based study: The Tromso Study. *Am J Epidemiol* 2001;154:236-44.

8. Forsdahl SH, Singh K, Solberg S, Jacobsen BK. Risk factors for abdominal aortic aneurysms: a 7-year prospective study: the Tromso Study, 1994-2001. *Circulation* 2009;119:2202-8.
9. Little J, Higgins JP, Ioannidis JP, Moher D, Gagnon F, von Elm E, et al. STrengthening the REporting of Genetic Association Studies (STREGA): an extension of the STROBE statement. *PLoS Med* 2009;6:e22.
10. Badger SA, Soong CV, O'Donnell ME, Mercer C, Young IS, Hughes AE. C-reactive protein (CRP) elevation in patients with abdominal aortic aneurysm is independent of the most important CRP genetic polymorphism. *J Vasc Surg* 2009;49:178-84.
11. Saratzis A, Bown M, Wild B, Sayers RD, Nightingale P, Smith J, et al. C-reactive protein polymorphism rs3091244 is associated with abdominal aortic aneurysm. *J Vasc Surg* 2014;60:1332-9.
12. Ioannidis JP, Boffetta P, Little J, O'Brien TR, Uitterlinden AG, Vineis P, et al. Assessment of cumulative evidence on genetic associations: interim guidelines. *Int J Epidemiol* 2008;37:120-32.
13. Ramsbottom D, Fitzgerald P, Grace PA, McAnena O, Burke P, Collins P, et al. Biochemical and molecular genetic studies of abdominal aortic aneurysm in an Irish population. *Eur J Vasc Surg* 1994;8:716-22.
14. Powell JT, MacSweeney ST, Greenhalgh RM, Turner RJ, Henney AM. Interaction between fibrillin genotype and blood pressure and the development of aneurysmal disease. *Ann N Y Acad Sci* 1996;800:198-207.

15. Ramsbottom D, O'Neill A, Sexton DM, Gafoor RA, Bouchier-Hayes D, Croke DT. The cholesteryl ester transfer protein (CETP) locus as a candidate gene in abdominal aortic aneurysm. *Clin Genet* 1997;51:241-5.
16. Hamano K, Ohishi M, Ueda M, Fujioka K, Katoh T, Zempo N, et al. Deletion polymorphism in the gene for angiotensin-converting enzyme is not a risk factor predisposing to abdominal aortic aneurysm. *Eur J Vasc Endovasc* 1999;18:158-61.
17. Wang X, Tromp G, Cole CW, Verloes A, Sakalihasan N, Yoon S, et al. Analysis of coding sequences for tissue inhibitor of metalloproteinases 1 (TIMP1) and 2 (TIMP2) in patients with aneurysms. *Matrix Biol* 1999;18:121-4.
18. Yoon S, Tromp G, Vongpunsawad S, Ronkainen A, Juvonen T, Kuivaniemi H. Genetic analysis of MMP3, MMP9, and PAI-1 in Finnish patients with abdominal aortic or intracranial aneurysms. *Biochem Bioph Res Com* 1999;265:563-8.
19. Kotani K, Shimomura T, Murakami F, Ikawa S, Kanaoka Y, Ohgi S, et al. Allele frequency of human endothelial nitric oxide synthase gene polymorphism in abdominal aortic aneurysm. *Intern Med* 2000;39:537-9.
20. Rossaak J, van Rij A, Jones G, Harris E. Association of the 4G/5G polymorphism in the promoter region of plasminogen activator inhibitor-1 with abdominal aortic aneurysms. *J Vasc Surg* 2000;31:1026-32.
21. Pola R, Gaetani E, Santoliquido A, Gerardino L, Cattani P, Serricchio M, et al. Abdominal aortic aneurysm in normotensive patients: association with angiotensin-converting enzyme gene polymorphism. *Eur J Vasc Endovasc* 2001;21:445-9.

22. Rasmussen T, Hallett J, Schulte S, Harmsen W, O'Fallon W, Weyand C. Genetic similarity in inflammatory and degenerative abdominal aortic aneurysms: A study of human leukocyte antigen class II disease risk genes. *J Vasc Surg* 2001;34:84-8.
23. Schillinger M, Exner M, Mlekusch W, Domanovits H, Huber K, Mannhalter C, et al. Heme oxygenase-1 gene promoter polymorphism is associated with abdominal aortic aneurysm. *Thromb Res* 2002;106:131-6.
24. Unno N, Nakamura T, Mitsuoka H, Uchiyama T, Yamamoto N, Saito T, et al. Association of a G994 -->T missense mutation in the plasma platelet-activating factor acetylhydrolase gene with risk of abdominal aortic aneurysm in Japanese. *Ann Surg* 2002;235:297-302.
25. Bown MJ, Burton PR, Horsburgh T, Nicholson ML, Bell PR, Sayers RD. The role of cytokine gene polymorphisms in the pathogenesis of abdominal aortic aneurysms: a case-control study. *J Vasc Surg* 2003;37:999-1005.
26. Jones G, Phillips V, Harris E, Rossaak J, van Rij A. Functional matrix metalloproteinase-9 polymorphism (C-1562T) associated with abdominal aortic aneurysm. *J Vasc Surg* 2003;38:1363-7.
27. Sugimoto T, Sada M, Miyamoto T, Yao H. Genetic analysis on HLA loci in Japanese patients with abdominal aortic aneurysm. *Eur J Vasc Endovasc* 2003;26:215-8.
28. Ghilardi G, Biondi ML, Battaglioli L, Zambon A, Guagnellini E, Scorza R. Genetic risk factor characterizes abdominal aortic aneurysm from arterial occlusive disease in human beings: CCR5 Delta 32 deletion. *J Vasc Surg* 2004;40:995-1000.

29. Massart F, Marini F, Menegato A, Del Monte F, Nuti M, Butitta F, et al. Allelic genes involved in artery compliance and susceptibility to sporadic abdominal aortic aneurysm. *J Steroid Biochem Mol Biol* 2004;92:413-8.
30. Fatini C, Pratesi G, Sofi F, Gensini F, Sticchi E, Lari B, et al. ACE DD genotype: A predisposing factor for abdominal aortic aneurysm. *Eur J Vasc Endovasc* 2005;29:227-32.
31. Fatini C, Sofi F, Sticchi E, Bolli P, Sestini I, Falciani M, et al. eNOS G894T polymorphism as a mild predisposing factor for abdominal aortic aneurysm. *J Vasc Surg* 2005;42:415-9.
32. Jones G, Harris E, Phillips L, van Rij A. The methylenetetrahydrofolate reductase C677T polymorphism does not associate with susceptibility to abdominal aortic aneurysm. *Eur J Vasc Endovasc* 2005;30:137-42.
33. Ogata T, Shibamura H, Tromp G, Sinha M, Goddard KA, Sakalihasan N, et al. Genetic analysis of polymorphisms in biologically relevant candidate genes in patients with abdominal aortic aneurysms. *J Vasc Surg* 2005;41:1036-42.
34. Schulz V, Hendig D, Schillinger M, Exner M, Domanovits H, Raith M, et al. Analysis of sequence variations in the ABCC6 gene among patients with abdominal aortic aneurysm and pseudoxanthoma elasticum. *J Vasc Res* 2005;42:424-32.
35. Strauss E, Waliszewski K, Pawlak AL. [The different genotypes of MTHFR 1298A>C and PON1 -108C>T polymorphisms confer the increased risk of the abdominal aortic aneurysm in the smoking and nonsmoking persons]. *Przegląd Lekarski* 2005;62:1023-30.

36. Ferrara F, Novo S, Grimaudo S, Raimondi F, Meli F, Amato C, et al. Methylenetetrahydrofolate reductase mutation in subjects with abdominal aortic aneurysm subdivided for age. *Clin Hemorheol Micro* 2006;34:421-6.
37. Hinterseher I, Bergert H, Kuhlisch E, Bloomenthal A, Pilarsky C, Ockert D, et al. Matrix metalloproteinase 2 polymorphisms in a caucasian population with abdominal aortic aneurysm. *J Surg Res* 2006;133:121-8.
38. Ogata T, Gregoire L, Goddard KAB, Skunca M, Tromp G, Lancaster WD, et al. Evidence for association between the HLA-DQA locus and abdominal aortic aneurysms in the Belgian population: A case control study. *BMC Med Genet* 2006;7.
39. Armani C, Curcio M, Barsotti MC, Santoni T, Di Stefano R, Dell'omodarme M, et al. Polymorphic analysis of the matrix metalloproteinase-9 gene and susceptibility to sporadic abdominal aortic aneurysm. *Biomed Pharmacother* 2007;61:268-71.
40. Badger SA, Soong CV, O'Donnell ME, Middleton D. The role of human leukocyte antigen genes in the formation of abdominal aortic aneurysms. *J Vasc Surg* 2007;45:475-80.
41. Bown MJ, Lloyd GM, Sandford RM, Thompson JR, London NJ, Samani NJ, et al. The interleukin-10-1082 'A' allele and abdominal aortic aneurysms. *J Vasc Surg* 2007;46:687-93.
42. Deguara J, Burnand KG, Berg J, Green P, Lewis CM, Chinien G, et al. An increased frequency of the 5A allele in the promoter region of the MMP3 gene is associated with abdominal aortic aneurysms. *Hum Mol Genet* 2007;16:3002-7.

43. Golledge J, Muller J, Shephard N, Clancy P, Smallwood L, Moran C, et al. Association between osteopontin and human abdominal aortic aneurysm. *Arterioscler Thromb Vasc* 2007;27:655-60.
44. Hinterseher I, Krex D, Kuhlisch E, Schmidt KG, Pilarsky C, Schneiders W, et al. Tissue inhibitor of metalloproteinase-1 (TIMP-1) polymorphisms in a caucasian population with abdominal aortic aneurysm. *World J Surg* 2007;31:2248-54.
45. Peeters AC, van Landeghem BA, Graafsma SJ, Kranendonk SE, Hermus AR, Blom HJ, et al. Low vitamin B6, and not plasma homocysteine concentration, as risk factor for abdominal aortic aneurysm: a retrospective case-control study. *J Vasc Surg* 2007;45:701-5.
46. Waliszewski K, Slomski R, Oszkinis G, Majewski W, Pawlaczyk K. Increased frequency of ace gene DD genotype in normotensive male smokers with abdominal aortic aneurysm. *Pol Przegl Chir* 2007;79:663-73.
47. Bown MJ, Braund PS, Thompson J, London NJM, Samani NJ, Sayers RD. Association Between the Coronary Artery Disease Risk Locus on Chromosome 9p21.3 and Abdominal Aortic Aneurysm. *Circ Cardiovasc Genet* 2008;1:39-42.
48. Giusti B, Saracini C, Bolli P, Magi A, Sestini I, Sticchi E, et al. Genetic analysis of 56 polymorphisms in 17 genes involved in methionine metabolism in patients with abdominal aortic aneurysm. *J Med Genet* 2008;45:721-30.
49. Gotting C, Prante C, Schillinger M, Exner M, Domanovits H, Raith M, et al. Xylosyltransferase I variants and their impact on abdominal aortic aneurysms. *Clinica Chimica Acta* 2008;391:41-5.

50. Helgadottir A, Gretarsdottir S, Thorleifsson G, Holm H, Patel RS, Gudnason T, et al. Apolipoprotein(a) Genetic Sequence Variants Associated With Systemic Atherosclerosis and Coronary Atherosclerotic Burden But Not With Venous Thromboembolism. *J Am Coll Cardiol* 2012;60:722-9.
51. Hinterseher I, Krex D, Kuhlisch E, Pilarsky C, Schneiders W, Saeger HD, et al. [Analysis of tissue inhibitor of metalloproteinase-2 gene polymorphisms in a caucasian population with abdominal aortic aneurysms]. *Zentralbl Chir* 2008;133:332-7.
52. Jones GT, Thompson AR, van Bockxmeer FM, Hafez H, Cooper JA, Golledge J, et al. Angiotensin II type 1 receptor 1166C polymorphism is associated with abdominal aortic aneurysm in three independent cohorts. *Arterioscler Thromb Vasc* 2008;28:764-70.
53. Smallwood L, Allcock R, van Bockxmeer F, Warrington N, Palmer LJ, Iacopetta B, et al. Polymorphisms of the interleukin-6 gene promoter and abdominal aortic aneurysm. *Eur J Vasc Endovasc* 2008;35:31-6.
54. Smallwood L, Allcock R, van Bockxmeer F, Warrington N, Palmer LJ, Iacopetta B, et al. Polymorphisms of the matrix metalloproteinase 9 gene and abdominal aortic aneurysm. *Br J Surg* 2008;95:1239-44.
55. Badger SA, Soong CV, Young IS, McGinty A, Mercer C, Hughes AE. The influence of COX-2 single nucleotide polymorphisms on abdominal aortic aneurysm development and the associated inflammation. *Angiology* 2009;60:576-81.

56. Elmore JR, Obmann MA, Kuivaniemi H, Tromp G, Gerhard GS, Franklin DP, et al. Identification of a genetic variant associated with abdominal aortic aneurysms on chromosome 3p12.3 by genome wide association. *J Vasc Surg* 2009;49:1525-31.
57. Elmore JR, Obmann MA, Kuivaniemi H, Tromp G, Gerhard GS, Franklin DP, et al. Regarding "Identification of a genetic variant associated with abdominal aortic aneurysms on chromosome 3p12.3 by genome wide association" Reply. *J Vasc Surg* 2009;50:1247.
58. Golledge J, Clancy P, Jones GT, Cooper M, Palmer LJ, van Rij AM, et al. Possible association between genetic polymorphisms in transforming growth factor beta receptors, serum transforming growth factor beta 1 concentration and abdominal aortic aneurysm. *Br J Surg* 2009;96:628-32.
59. Jones GT, van Rij AM. Regarding "Identification of a genetic variant associated with abdominal aortic aneurysms on chromosome 3p12.3 by genome wide association". *J Vasc Surg* 2009;50:1246-7.
60. Korcz A, Mikolajczyk-Stecyna J, Gabriel M, Zowczak-Drabarczyk M, Pawlaczyk K, Kalafirov M, et al. Angiotensin-Converting Enzyme (ACE, I/D) Gene Polymorphism and Susceptibility to Abdominal Aortic Aneurysm or Aortoiliac Occlusive Disease. *J Surg Res* 2009;153:76-82.
61. Lucarini L, Sticchi E, Sofi F, Pratesi G, Pratesi C, Pulli R, et al. ACE and TGFB1 genes interact in influencing the susceptibility to abdominal aortic aneurysm. *Atherosclerosis* 2009;202:205-10.
62. Sandford B, Bown M, London N, Sayers R. The role of the CCR5 Delta32 polymorphism in abdominal aortic aneurysms. *Int J Immunogenet* 2009;36:199-205.

63. Smallwood L, Warrington N, Allcock R, van Bockxmeer F, Palmer LJ, Iacopetta B, et al. Matrix Metalloproteinase-2 Gene Variants and Abdominal Aortic Aneurysm. *Eur J Vasc Endovasc* 2009;38:169-71.
64. Thompson AR, Golledge J, Cooper JA, Hafez H, Norman PE, Humphries SE. Sequence variant on 9p21 is associated with the presence of abdominal aortic aneurysm disease but does not have an impact on aneurysmal expansion. *Eur J Hum Genet* 2009;17:391-4.
65. Atli FH, Manduz S, Katrancioğlu N, Ozum U, Disli OM, Atahan E, et al. eNOS G894T polymorphism and abdominal aortic aneurysms. *Angiology* 2010;61:125-30.
66. Baas AF, Medic J, van't Slot R, de Vries JP, van Sambeek MR, Geelkerken BH, et al. Association study of single nucleotide polymorphisms on chromosome 19q13 with abdominal aortic aneurysm. *Angiology* 2010;61:243-7.
67. Baas AF, Medic J, van't Slot R, de Kovel CG, Zhernakova A, Geelkerken RH, et al. Association of the TGF-beta receptor genes with abdominal aortic aneurysm. *Eur J Hum Genet* 2010;18:240-4.
68. Baas AF, Medic J, van't Slot R, de Vries JP, van Sambeek MR, Verhoeven EL, et al. The intracranial aneurysm susceptibility genes HSPG2 and CSPG2 are not associated with abdominal aortic aneurysm. *Angiology* 2010;61:238-42.
69. Badger SA, Soong CV, O'Donnell ME, Sharif MA, Makar RR, Hughes AE. Common polymorphisms of Fibulin-5 and the risk of abdominal aortic aneurysm development. *Vasc Med* 2010;15:113-7.

70. Biros E, Cooper M, Palmer LJ, Walker PJ, Norman PE, Golledge J. Association of an allele on chromosome 9 and abdominal aortic aneurysm. *Atherosclerosis* 2010;212:539-42.
71. Golledge J, Biros E, Cooper M, Warrington N, Palmer LJ, Norman PE. Apolipoprotein E genotype is associated with serum C-reactive protein but not abdominal aortic aneurysm. *Atherosclerosis* 2010;209:487-91.
72. Gretarsdottir S, Baas AF, Thorleifsson G, Holm H, den Heijer M, de Vries JP, et al. Genome-wide association study identifies a sequence variant within the DAB2IP gene conferring susceptibility to abdominal aortic aneurysm. *Nat Genet* 2010;42:692-7.
73. Moran CS, Clancy P, Biros E, Blanco-Martin B, McCaskie P, Palmer LJ, et al. Association of PPARgamma allelic variation, osteoprotegerin and abdominal aortic aneurysm. *Clin Endocrinol (Oxf)* 2010;72:128-32.
74. Obukofe B, Sayers RD, Thompson J, Sandford RM, London NJM, Samani NJ, et al. Is there an association between angiotensin converting enzyme (ACE) genotypes and abdominal aortic aneurysm?. *Eur J Vasc Endovasc* 2010;40:457-60.
75. Thompson AR, Cooper JA, Jones GT, Drenos F, van Bockxmeer FM, Biros E, et al. Assessment of the association between genetic polymorphisms in transforming growth factor beta, and its binding protein (LTBP), and the presence, and expansion, of Abdominal Aortic Aneurysm. *Atherosclerosis* 2010;209:367-73.
76. Biros E, Norman PE, Walker PJ, Nataatmadja M, West M, Golledge J. A single nucleotide polymorphism in exon 3 of the kallikrein 1 gene is associated with large but not small abdominal aortic aneurysm. *Atherosclerosis* 2011;217:452-7.

77. Biros E, Norman PE, Jones GT, van Rij AM, Yu G, Moxon JV, et al. Meta-analysis of the association between single nucleotide polymorphisms in TGF-receptor genes and abdominal aortic aneurysm. *Atherosclerosis* 2011;219:218-23.
78. Bown MJ, Jones GT, Harrison SC, Wright BJ, Bumpstead S, Baas AF, et al. Abdominal Aortic Aneurysm Is Associated with a Variant in Low-Density Lipoprotein Receptor-Related Protein 1. *Am J Hum Genet* 2011;89:619-27.
79. Bradley DT, Badger SA, Bown MJ, Sayers RD, Hughes AE. Coding polymorphisms in the genes of the alternative complement pathway and abdominal aortic aneurysm. *Int J Immunogenet* 2011;38:243-8.
80. Hinterseher I, Erdman R, Donoso LA, Vrabec TR, Schworer CM, Lillvis JH, et al. Role of complement cascade in abdominal aortic aneurysms. *Arteriosc Thromb Vas* 2011;31:1653-60.
81. Katrancioglu N, Berkan O, Manduz S, Atli FH, Yilmaz MB. ENOS G894T polymorphism and abdominal aortic aneurysms. *Angiology* 2010;61:227-8.
82. Lillvis JH, Kyo Y, Tromp G, Lenk GM, Li M, Lu Q, et al. Analysis of positional candidate genes in the AAA1 susceptibility locus for abdominal aortic aneurysms on chromosome 19. *BMC Med Genet* 2011;12.
83. Roberts RL, Van Rij AM, Phillips LV, Young S, McCormick SP, Merriman TR, et al. Interaction of the inflammasome genes CARD8 and NLRP3 in abdominal aortic aneurysms. *Atherosclerosis* 2011;218:123-6.
84. Bisioendial RJ, Tanck MW, Golledge J, Broekhuizen LN, Legemate DA, Stroes ES, et al. The association between the gene encoding 5-lipoxygenase activating protein and abdominal aortic aneurysms. *Atherosclerosis* 2012;220:425-8.

85. Duellman T, Warren CL, Peissig P, Wynn M, Yang J. Matrix Metalloproteinase-9 Genotype as a Potential Genetic Marker for Abdominal Aortic Aneurysm. *Circ Cardiovasc Genet* 2012;5:529-37.
86. Korcz A, Zakerska O, Pawlaczyk K, Szypura-Ratajczak D, Molińska-Glura M, Oszkinis G, et al. Analiza polimorfizmu genu metaloproteinazy 13 (-77A> G) w polskiej populacji pacjentów z tętniakami aorty brzusznej iz niedrożnością aortalno-biodrową. *Acta Angiologica* 2012;18:148-56.
87. Oszejka K, Wroński K, Janiszewska G, Bieńkiewicz M, Bartkowiak J, Szemraj J. The study of t-PA, u-PA and PAI-1 genes polymorphisms in patients with abdominal aortic aneurysm. *Mol Biol Rep* 2014;41:2859-64.
88. Saracini C, Bolli P, Sticchi E, Pratesi G, Pulli R, Sofi F, et al. Polymorphisms of genes involved in extracellular matrix remodeling and abdominal aortic aneurysm. *J Vasc Surg* 2012;55:171-9.
89. Antoniou GA, Lazarides MK, Patera S, Antoniou SA, Giannoukas AD, Georgiadis GS, et al. Assessment of insertion/deletion polymorphism of the angiotensin-converting enzyme gene in abdominal aortic aneurysm and inguinal hernia. *Vascular* 2013;21:1-5.
90. Bradley DT, Hughes AE, Badger SA, Jones GT, Harrison SC, Wright BJ, et al. A variant in LDLR is associated with abdominal aortic aneurysm. *Circ Cardiovasc Genet* 2013;6:498-504.
91. Galora S, Saracini C, Palombella AM, Pratesi G, Pulli R, Pratesi C, et al. Low-density lipoprotein receptor-related protein 5 gene polymorphisms and genetic susceptibility to abdominal aortic aneurysm. *J Vasc Surg* 2013;58:1062-68.

92. Gregorek AC, Gornik KC, Polancec DS, Dabelic S. GT microsatellite repeats in the heme oxygenase-1 gene promoter associated with abdominal aortic aneurysm in Croatian patients. *Biochem Genet* 2013;51:482-92.
93. Harrison SC, Smith AJP, Jones GT, Swerdlow DI, Rampuri R, Bown MJ, et al. Interleukin 6 Receptor Pathways in Abdominal Aortic Aneurysm. *Eur Heart J* 2012; 34:3707-16.
94. Jones GT, Bown MJ, Gretarsdottir S, Romaine SPR, Helgadottir A, Yu G, et al. A sequence variant associated with Sortilin-1 (SORT1) on 1p13.3 is independently associated with Abdominal Aortic Aneurysm. *Hum Mol Genet* 2013;22:2941-7.
95. Tragante V, Doevendans PA, Nathoe HM, van der Graaf Y, Spiering W, Algra A, et al. The impact of susceptibility loci for coronary artery disease on other vascular domains and recurrence risk. *Eur Heart J* 2013;34:2896-904.
96. van 't Hof FN, Ruigrok YM, Baas AF, Kiemeney LA, Vermeulen SH, Uitterlinden AG, et al. Impact of inherited genetic variants associated with lipid profile, hypertension, and coronary artery disease on the risk of intracranial and abdominal aortic aneurysms. *Circ Cardiovasc Genet* 2013;6:264-70.
97. Wong YY, Golledge J, Flicker L, McCaul KA, Hankey GJ, van Bockxmeer FM, et al. Plasma total homocysteine is associated with abdominal aortic aneurysm and aortic diameter in older men. *J Vasc Surg* 2013;58:364-70.
98. Bridge KI, Macrae F, Bailey MA, Johnson A, Philippou H, Scott DJA, et al. The alpha-2-antiplasmin Arg407Lys polymorphism is associated with Abdominal Aortic Aneurysm. *Thromb Res* 2014;134:723-8.

99. Cao H, Hu X, Zhang Q, Li J, Liu B, Wang J, et al. Hyperhomocysteinaemia, low folate concentrations and MTHFR C677T mutation in abdominal aortic aneurysm. *Vasa* 2014;43:181-8.
100. Duellman T, Warren CL, Peissig P, Wynn M, Yang J. Matrix metalloproteinase-9 genotype as a potential genetic marker for abdominal aortic aneurysm. *Circ Cardiovasc Genet* 2012;5:529-37.
101. Galora S, Saracini C, Pratesi G, Sticchi E, Pulli R, Pratesi C, et al. Association of rs1466535 LRP1 but not rs3019885 SLC30A8 and rs6674171 TDRD10 gene polymorphisms with abdominal aortic aneurysm in Italian patients. *J Vasc Surg* 2014.
102. Li Y, Yang C, Ma G, Cui L, Gu X, Chen Y, et al. Analysis of ADAM17 polymorphisms and susceptibility to sporadic abdominal aortic aneurysm. *Cell Physiol Biochem* 2014;33:1426-38.
103. Mikolajczyk-Stecyna J, Korcz A, Gabriel M, Pawlaczyk K, Oszkinis G, Slomski R. Risk factors in abdominal aortic aneurysm and in Polish population aortoiliac occlusive disease and differences between them. *Sci Rep* 2013;3:3528.
104. Oszejca K, Wroński K, Janiszewska G, Bieńkiewicz M, Panek M, Bartkowiak J, et al. Association analysis of genetic polymorphisms of factor V, factor VII and fibrinogen β chain genes with human abdominal aortic aneurysm. *Exp Ther Med* 2012;4:514-8.
105. Saratzis A, Bown MJ, Wild B, Nightingale P, Smith J, Johnson C, et al. Association between seven single nucleotide polymorphisms involved in inflammation and proteolysis and abdominal aortic aneurysm. *J Vasc Surg* 2014.

106. Strauss E, Waliszewski K, Oszkinis G, Staniszewski R. Polymorphisms of genes involved in the hypoxia signaling pathway and the development of abdominal aortic aneurysms or large-artery atherosclerosis. *J Vasc Surg* 2014.
107. Wei Y, Xiong J, Zuo S, Chen F, Chen D, Wu T, et al. Association of polymorphisms on chromosome 9p21. 3 region with increased susceptibility of abdominal aortic aneurysm in a Chinese Han population. *J Vasc Surg* 2014;59:879-85.
108. Helgadóttir A, Thorleifsson G, Magnusson KP, Gretarsdóttir S, Steinthorsdóttir V, Manolescu A, et al. The same sequence variant on 9p21 associates with myocardial infarction, abdominal aortic aneurysm and intracranial aneurysm. *Nat Genet* 2008;40:217-24.
109. Etique N, Verzeaux L, Dedieu S, Emonard H. LRP-1: a checkpoint for the extracellular matrix proteolysis. *BioMed Res Int* 2013;2013:152163.
110. Kamstrup PR. Lipoprotein (a) and ischemic heart disease—a causal association? A review. *Atherosclerosis* 2010;211:15-23.
111. Elliott P, Chambers JC, Zhang W, Clarke R, Hopewell JC, Peden JF, et al. Genetic Loci associated with C-reactive protein levels and risk of coronary heart disease. *JAMA* 2009;302:37-48.
112. Dehghan A, Dupuis J, Barbalic M, Bis JC, Eiriksdóttir G, Lu C, et al. Meta-analysis of genome-wide association studies in >80 000 subjects identifies multiple loci for C-reactive protein levels. *Circulation* 2011;123:731-8.
113. IL6R Genetics Consortium Emerging Risk Factors, Collaboration, Sarwar N, Butterworth AS, Freitag DF, Gregson J, Willeit P, et al. Interleukin-6 receptor

pathways in coronary heart disease: a collaborative meta-analysis of 82 studies. Lancet 2012;379:1205-13.

114. Coronary Artery Disease (C4D) Genetics Consortium. A genome-wide association study in Europeans and South Asians identifies five new loci for coronary artery disease. Nat Genet 2011;43:339-44.

115. Newman KM, Ogata Y, Malon AM, Irizarry E, Gandhi RH, Nagase H, et al. Identification of matrix metalloproteinases 3 (stromelysin-1) and 9 (gelatinase B) in abdominal aortic aneurysm. Arterioscler Thromb 1994;14:1315-20.

116. Morris DR, Biros E, Cronin O, Kuivaniemi H, Golledge J. The association of genetic variants of matrix metalloproteinases with abdominal aortic aneurysm: a systematic review and meta-analysis. Heart 2014;100:295-302.

117. Wang W, Zee RL, Morris BJ. Association of angiotensin II type 1 receptor gene polymorphism with essential hypertension. Clin Genet 1997;51:31-4.

Tables

Table 1. Databases and time periods searched

Database	Date Range
Medline	1946 - August Week 1 2014
Embase	1980 – 2014 Week 33
Web of Science	All years – 25 August 2014

Table 2. Study Quality

First Author	Year	Cases (n)	Controls (n)	Ethnicity reported	Controlled for ethnicity by logistic regression	Tested or untested controls	Genotyping rate reported	HWE Reported	Multiple tests	Corr. for Multiple Testing	Study Type	Reported Association
Ramsbottom ¹³	1994	82	79	Yes	No	Tested	No	No	Yes	No	Cand	No
Powell ¹⁴	1996	232	245	Yes	No	Untested	No	No	Yes	No	Cand	No
Ramsbottom ¹⁵	1997	85	34	No	No	Tested	No	No	Yes	No	Cand	No
Hamano ¹⁶	1999	125	153	No	No	Tested	No	Yes	No	NA	Cand	No
Wang ¹⁷	1999	84	51	No	No	NR	No	No	Yes	No	Cand	Yes
Yoon ¹⁸	1999	47	174	No	No	Untested	Yes	Yes	Yes	Yes	Cand	No
Kotani ¹⁹	2000	58	410	No	No	Untested	No	Yes	Yes	No	Cand	No
Rossaak ²⁰	2000	190	163	Yes	No	Untested	No	No	Yes	Yes	Cand	Yes
Pola ²¹	2001	124	112	Yes	No	Tested	No	Yes	Yes	No	Cand	No
Rasmussen ²²	2001	102	118	Yes	No	Untested	No	No	Yes	No	Cand	Yes
Schillinger ²³	2002	70	61	No	No	Tested	No	No	Yes	No	Cand	Yes
Unno ²⁴	2002	131	106	No	No	Untested	No	Yes	Yes	No	Cand	Yes
Bown ²⁵	2003	100	100	Yes	No	Mixed	No	Yes	Yes	No	Cand	Yes
Jones ²⁶	2003	414	203	No	No	Untested	No	Yes	Yes	No	Cand	Yes
Sugimoto ²⁷	2003	49	237	No	No	Untested	No	No	Yes	No	Cand	Yes
Ghilardi ²⁸	2004	70	172	No	No	Tested	No	Yes	Yes	No	Cand	Yes
Massart ²⁹	2004	99	225	Yes	No	NR	No	Yes	Yes	No	Cand	Yes
Fatini ³⁰	2005	250	250	Yes	No	Tested	No	Yes	Yes	No	Cand	Yes
Fatini ³¹	2005	250	250	Yes	No	Tested	No	Yes	Yes	No	Cand	Yes
Jones ³²	2005	428	282	No	No	Untested	No	Yes	Yes	No	Cand	No
Ogata ³³	2005	387	425	Yes	Yes	Untested	No	Yes	Yes	No	Cand	Yes
Schulz ³⁴	2005	133	910	Yes	No	Untested	No	No	Yes	No	Cand	No

First Author	Year	Cases (n)	Controls (n)	Ethnicity reported	Controlled for ethnicity by logistic regression	Tested or untested controls	Genotyping rate reported	HWE Reported	Multiple tests	Corr. for Multiple Testing	Study Type	Reported Association
Strauss ³⁵	2005	106	97	No	No	Untested	No	Yes	Yes	No	Cand	Yes
Ferrara ³⁶	2006	88	44	No	No	NR	No	No	Yes	Yes	Cand	Yes
Hinterseher ³⁷	2006	51	48	Yes	No	Untested	No	Yes	Yes	No	Cand	No
Ogata ³⁸	2006	387	426	Yes	No	Untested	No	No	Yes	Yes	Cand	Yes
Armani ³⁹	2007	146	156	No	No	Untested	No	No	Yes	No	Cand	No
Badger ⁴⁰	2007	241	1000	No	No	Untested	No	Yes	Yes	Yes	Cand	No
Bown ⁴¹	2007	389	404	Partial	No	Tested	No	Yes	Yes	No	Cand	Yes
Deguarra ⁴²	2007	405	405	Yes	No	Untested	No	No	Yes	No	Cand	Yes
Golledge ⁴³	2007	689	3538	No	No	Tested	Yes	Yes	Yes	Yes	Cand	No
Hinterseher ⁴⁴	2007	146	133	Yes	No	Untested	No	Yes	Yes	No	Cand	No
Peeters ⁴⁵	2007	88	88	No	No	Tested	No	No	No	NA	Cand	No
Waliszewski ⁴⁶	2007	112	50	Yes	No	Untested	No	No	Yes	No	Cand	Yes
Bown ⁴⁷	2008	899	815	Yes	No	Tested	No	Yes	Yes	No	Cand	Yes
Giusti ⁴⁸	2008	423	423	No	No	Tested	No	Yes	Yes	Yes	Cand	Yes
Gotting ⁴⁹	2008	129	129	No	No	Untested	No	Yes	Yes	Yes	Cand	Yes
Helgadottir ⁵⁰	2008	2836	16732	Partial	No	Mixed	No	Yes	Yes	No	Cand	Yes
Hinterseher ⁵¹	2008	50	41	Partial	No	Untested	No	Yes	Yes	No	Cand	Yes
Jones ⁵²	2008	1226	1723	Partial	No	Mixed	No	Yes	Yes	No	Cand	Yes
Smallwood ⁵³	2008	677	656	No	No	Tested	Yes	Yes	Yes	No	Cand	Yes
Smallwood ⁵⁴	2008	678	659	No	No	Tested	Yes	Yes	Yes	No	Cand	No
Badger ⁵⁵	2009	230	279	No	No	Tested	No	No	Yes	No	Cand	No
Badger ¹⁰	2009	248	400	No	No	Tested	No	No	No	NA	Cand	No
Elmore ^{56,57}	2009	950	1146	No	No	Mixed	No	No	Yes	No	GWAS	Yes
Golledge ⁵⁸	2009	1294	1460	No	No	Tested	Yes	Yes	Yes	Yes	Cand	No
Jones ⁵⁹	2009	567	552	Yes	NR	Tested	No	No	Yes	No	Cand	No

First Author	Year	Cases (n)	Controls (n)	Ethnicity reported	Controlled for ethnicity by logistic regression	Tested or untested controls	Genotyping rate reported	HWE Reported	Multiple tests	Corr. for Multiple Testing	Study Type	Reported Association
Korcz ⁶⁰	2009	133	152	Yes	No	Tested	No	Yes	Yes	No	Cand	No
Lucarini ⁶¹	2009	201	252	No	No	Tested	No	Yes	Yes	No	Cand	Yes
Sandford ⁶²	2009	285	273	Yes	No	Tested	No	No	Yes	No	Cand	No
Smallwood ⁶³	2009	678	659	No	No	Tested	Yes	Yes	Yes	Yes	Cand	Yes
Thompson ⁶⁴	2009	741	1366	No	No	Partial	No	No	Yes	No	Cand	Yes
Atli ⁶⁵	2010	61	62	No	No	Tested	No	No	Yes	No	Cand	Yes
Baas ⁶⁶	2010	736	1024	No	No	Mixed	Yes	Yes	Yes	Yes	Cand	No
Baas ⁶⁷	2010	736	1024	No	No	Mixed	Yes	Yes	Yes	Yes	Cand	Yes
Baas ⁶⁸	2010	736	1024	No	No	Mixed	Yes	Yes	Yes	Yes	Cand	No
Badger ⁶⁹	2010	230	278	Yes	No	Tested	No	Yes	Yes	No	Cand	No
Biros ⁷⁰	2010	513	2858	No	No	Tested	No	No	Yes	No	Cand	Yes
Golledge ⁷¹	2010	640	1071	No	No	Tested	Yes	Yes	Yes	Yes	Cand	No
Gretarsdottir ⁷²	2010	4559	37954	Partial	No	Mixed	No	No	Yes	Yes	GWAS	Yes
Moran ⁷³	2010	689	3538	No	No	Tested	Yes	Yes	Yes	No	Cand	Yes
Obukofe ⁷⁴	2010	1155	996	No	No	Tested	No	Yes	Yes	No	Cand	No
Thompson ⁷⁵	2010	1890	3785	Yes	No	Mixed	No	Yes	Yes	No	Cand	No
Biros ⁷⁶	2011	834	795	No	No	Tested	No	Yes	Yes	No	Cand	Yes
Biros ⁷⁷	2011	1904	2616	No	No	Mixed	No	Yes	Yes	Yes	Cand	No
Bown ⁷⁸	2011	6228	49182	No	Partial	Mixed	No	Yes	Yes	Yes	GWAS	Yes
Bradley ⁷⁹	2011	434	378	Yes	No	Tested	Yes	Yes	Yes	Yes	Cand	No
Hinterseher ⁸⁰	2011	874	899	No	No	NR	No	Yes	Yes	No	Cand	No
Katrancioglu ⁸¹	2011	100	138	No	No	Tested	No	No	Yes	No	Cand	Yes
Lillvis ⁸²	2011	394	419	No	No	Untested	No	Yes	Yes	Yes	Cand	Yes
Roberts ⁸³	2011	1238	731	Yes	No	Tested	Yes	Yes	Yes	No	Cand	Yes
Bisoendial ⁸⁴	2012	613	707	No	No	Tested	No	Yes	Yes	No	Cand	No

First Author	Year	Cases (n)	Controls (n)	Ethnicity reported	Controlled for ethnicity by logistic regression	Tested or untested controls	Genotyping rate reported	HWE Reported	Multiple tests	Corr. for Multiple Testing	Study Type	Reported Association
Duellman ⁸⁵	2012	178	178	No	No	Tested	No	Yes	Yes	No	Cand	Yes
Helgadottir ⁵⁰	2012	4261	33520	Partial	No	Mixed	No	Yes	Yes	Yes	Cand	Yes
Korcz ⁸⁶	2012	300	313	No	No	Tested	No	Yes	Yes	No	Cand	No
Oszajca ⁸⁷	2012	153	152	No	No	Untested	No	Yes	Yes	No	Cand	Yes
Saracini ⁸⁸	2012	423	423	No	No	Tested	No	Yes	Yes	Yes	Cand	Yes
Antoniou ⁸⁹	2013	65	89	Yes	No	Tested	No	No	No	NA	Cand	Yes
Bradley ⁹⁰	2013	5138	39273	Partial	Partial	Mixed	Yes	Yes	Yes	Yes	GWAS	Yes
Galora ⁹¹	2013	423	423	No	No	Tested	No	Yes	Yes	Yes	Cand	Yes
Gregorek ⁹²	2013	117	117	No	No	Tested	No	Yes	Yes	No	Cand	Yes
Harrison ⁹³	2013	4524	15710	Partial	No	Mixed	Yes	Yes	No	NA	Cand	Yes
Jones ⁹⁴	2013	7048	75976	Partial	No	Mixed	Yes	Yes	Yes	No	Cand	Yes
Tragante ⁹⁵	2013	651	2015	No	No	Untested	Yes	Yes	Yes	Yes	GRS	Yes
van 't Hof ⁹⁶	2013	859	2089	Yes	Yes	Untested	Yes	Yes	Yes	No	GRS	Yes
Wong ⁹⁷	2013	318	3930	No	No	Tested	No	Yes	Yes	No	Cand	No
Bridge ⁹⁸	2014	602	490	Yes	No	Tested	Yes	Yes	Yes	No	Cand	Yes
Cao ⁹⁹	2014	463	463	Yes	No	Tested	No	Yes	Yes	No	Cand	Yes
Duellman ¹⁰⁰	2014	141	168	No	No	Tested	Yes	Yes	Yes	No	Cand	Yes
Galora ¹⁰¹	2014	423	423	No	No	Untested	No	Yes	Yes	Yes	Cand	Yes
Li ¹⁰²	2014	316	306	No	No	Tested	No	Yes	Yes	No	Cand	Yes
Mikołajczyk-Stecyna ¹⁰³	2014	128	180	No	No	Tested	No	Yes	Yes	No	Cand	Yes
Oszajca ¹⁰⁴	2014	153	152	No	No	Untested	No	No	Yes	No	Cand	Yes
Saratzis ¹⁰⁵	2014	797	793	Partial	No	Tested	No	Yes	Yes	No	Cand	No
Saratzis ¹¹	2014	722	753	Partial	No	Tested	No	Yes	Yes	No	Cand	Yes
Strauss ¹⁰⁶	2014	518	541	No	No	Untested	No	Yes	Yes	No	Cand	Yes

First Author	Year	Cases (n)	Controls (n)	Ethnicity reported	Controlled for ethnicity by logistic regression	Tested or untested controls	Genotyping rate reported	HWE Reported	Multiple tests	Corr. for Multiple Testing	Study Type	Reported Association
Wei ¹⁰⁷	2014	155	310	Yes	No	Tested	Yes	Yes	Yes	No	Cand	Yes

NR, Not Reported; NA, Not Applicable; Cand, Candidate Gene Study; GRS, Genetic Risk Score; GWAS, Genome-wide Association

Study

Table 3. Summary of meta-analyses

Gene and Polymorphism	OR (95% CI)	p value	I ²	FE/RE
9p21 (<i>CDKN2BAS</i>) rs10757278 Heterozygous (A/G)	1.28 (1.16-1.40)	2.9x10 ⁻⁷	7.3%	FE
9p21 (<i>CDKN2BAS</i>) rs10757278 Homozygous (G/G)	1.55 (1.38-1.74)	1.2x10 ⁻¹³	20.1%	FE
<i>ACE</i> Insertion/Deletion (rs4646994) Heterozygous (D/I)	0.90 (0.72-1.12)	0.34	66.5%	RE
<i>ACE</i> Insertion/Deletion (rs4646994) Homozygous (I/I)	0.67 (0.50-0.90)	8.0x10 ⁻³	72.3%	RE
* <i>ACE</i> Insertion/Deletion (rs4646994) Heterozygous (D/I)	0.85 (0.71-1.02)	0.08	48.7%	RE
* <i>ACE</i> Insertion/Deletion (rs4646994) Homozygous (I/I)	0.67 (0.49-0.91)	9.5 x10 ⁻³	75.1%	RE
<i>AGTR1</i> rs5186 Heterozygous (A/C)	1.25 (1.00-1.55)	0.047	53.0%	RE
<i>AGTR1</i> rs5186 Homozygous (C/C)	1.44 (1.13-1.84)	3.5x10 ⁻³	0.6%	FE
<i>AGTR1</i> rs5186 Allelic Model (C Allele)	1.19 (1.05-1.34)	5.0x10 ⁻³	45.6%	RE
<i>APOA1</i> rs964184 Additive (G allele)	1.20 (1.05-1.37)	6.8x10 ⁻³	0.0%	FE
<i>APOB</i> rs1367117 Additive (A Allele)	1.09 (0.95-1.24)	0.24	50.1%	RE
<i>CCR5</i> rs333 Heterozygous (WT/Del32)	1.34 (0.36-4.94)	0.66	89.5%	RE
<i>CCR5</i> rs333 Homozygous (Del32/ Del32)	1.73 (0.21-14.09)	0.61	41.2%	RE
<i>CRP</i> rs3091244 Heterozygous (Group B)	1.58 (0.91-2.77)	0.11	87.4%	RE
<i>CRP</i> rs3091244 Homozygous (Group A)	2.16 (1.10-4.24)	0.03	84.0%	RE
* <i>CRP</i> rs3091244 Heterozygous (Group B)	1.25 (0.68-2.30)	0.46	82.5%	RE
* <i>CRP</i> rs3091244 Homozygous (Group A)	1.60 (0.79-3.22)	0.19	74.3%	RE
<i>ELN</i> rs2071307 Heterozygous (A/G)	0.73 (0.46-1.14)	0.16	73.5%	RE
<i>ELN</i> rs2071307 Homozygous (A/A)	0.90 (0.48-1.69)	0.74	77.9%	RE
<i>HMOX1</i> GT(n) Repeat Heterozygous (GT(≥25)/GT(<25))	1.21 (0.25-5.85)	0.81	91.7%	RE
<i>HMOX1</i> GT(n) Repeat Homozygous (GT(<25)/ GT(<25))	1.03 (0.14-7.58)	0.97	76.4%	RE
<i>IL10</i> rs1800896 Heterozygous (G/A)	1.26 (0.89-1.78)	0.20	31.1%	RE
<i>IL10</i> rs1800896 Homozygous (A/A)	1.59 (0.97-2.61)	0.06	53.7%	RE
<i>LRP1</i> rs1466535 Heterozygous (C/T)	0.93 (0.45-1.93)	0.84	57.5%	RE
<i>LRP1</i> rs1466535 Homozygous (C/C)	0.85 (0.22-3.32)	0.81	87.8%	RE

<i>LRP1</i> rs1466535 Additive (C Allele)	1.09 (1.00-1.19)	0.047	69.5%	RE
* <i>LRP1</i> rs1466535 Additive (C Allele)	1.15 (1.10-1.21)	2.5x10 ⁻¹⁰	0.0%	FE
<i>MMP2</i> rs243865 Heterozygous (C/T)	0.87 (0.60-1.24)	0.43	73.7%	RE
<i>MMP2</i> rs243865 Homozygous (T/T)	0.65 (0.46-0.93)	0.02	0%	FE
<i>MMP2</i> rs243865 Dominant (C/T & T/T)	0.82 (0.60-0.15)	0.25	70.9%	RE
<i>MMP3</i> rs3025058 Heterozygous (6A/5A)	0.84 (0.70-1.02)	0.08	0.0%	FE
<i>MMP3</i> rs3025058 Homozygous (6A/6A)	0.61 (0.49-0.76)	1.6x10 ⁻⁵	0.0%	FE
<i>MMP9</i> rs3918242 Heterozygous (C/T)	1.05 (0.92-1.18)	0.48	21.4%	FE
<i>MMP9</i> rs3918242 Homozygous (T/T)	0.81 (0.49-1.33)	0.41	35.4%	RE
* <i>MMP9</i> rs3918242 Heterozygous (C/T)	1.08 (0.91-1.27)	0.40	30.7%	RE
* <i>MMP9</i> rs3918242 Homozygous (T/T)	0.93 (0.61-1.42)	0.75	0%	FE
<i>MMP13</i> rs2252070 Heterozygous (A/G)	1.08 (0.90-1.29)	0.41	0.0%	FE
<i>MMP13</i> rs2252070 Homozygous (G/G)	1.16 (0.76-1.77)	0.48	50.7%	RE
<i>MTHFR1</i> rs1801133 Heterozygous (C/T)	1.06 (0.87-1.30)	0.57	68.6%	RE
<i>MTHFR1</i> rs1801133 Homozygous (T/T)	1.06 (0.81-1.40)	0.66	60.7%	RE
* <i>MTHFR1</i> rs1801133 Heterozygous (C/T)	1.07 (0.85-1.36)	0.56	72.3%	RE
* <i>MTHFR1</i> rs1801133 Homozygous (T/T)	0.96 (0.75-1.21)	0.70	37.1%	RE
<i>NOS3</i> rs1799983 Heterozygous (G/T)	1.14 (0.84-1.55)	0.39	29.9%	RE
<i>NOS3</i> rs1799983 Homozygous (T/T)	1.16 (0.52-2.58)	0.71	87.9%	RE
<i>PHACTR1</i> rs12526453 Additive (C allele)	1.02 (0.86-1.22)	0.80	63.7%	RE
<i>SERPINE1</i> rs1799889 Heterozygous (4G/5G)	0.90 (0.65-1.24)	0.51	0%	FE
<i>SERPINE1</i> rs1799889 Homozygous (5G/5G)	1.10 (0.72-1.69)	0.66	0%	FE
* <i>SERPINE1</i> rs1799889 Heterozygous (4G/5G)	0.92 (0.62-1.38)	0.62	0%	FE
* <i>SERPINE1</i> rs1799889 Homozygous (5G/5G)	1.19 (0.70-2.03)	0.52	0%	FE
<i>SORT1</i> rs599839 Additive (G Allele)	0.82 (0.77-0.86)	2.6x10 ⁻¹³	0.0%	FE
<i>TGFBR2</i> rs1036095 Heterozygous (C/G)	1.23 (0.83-1.81)	0.31	83.1%	RE
<i>TGFBR2</i> rs1036095 Homozygous (G/G)	1.69 (0.76-3.77)	0.20	88.5%	RE
<i>TGFBR2</i> rs764522 Heterozygous (C/G)	1.35 (0.91-1.99)	0.13	79.9%	RE
<i>TGFBR2</i> rs764522 Homozygous (G/G)	1.73 (0.78-3.85)	0.18	86.2%	RE

<i>TRIB1</i> rs2954029 Additive (A Allele)	1.06 (0.90-1.24)	0.45	59.0%	RE
--	------------------	------	-------	----

* Sensitivity Analysis conducted with exclusion of studies because of deviation from HWE or an extreme outlying result

OR, Odds Ratio; RE, Random Effects Meta-analysis; FE, Fixed Effects Meta-analysis

Table 4. Assessment of Evidence of Association

Gene	Polymorphism	Amount of Evidence	Replication	Protection from Bias	Overall Assessment
3p21	rs7635818	A	C	C	Weak
9p21	rs10757278	A	A	A	Strong
<i>ACE</i>	rs4646994	A	A	C	Weak
<i>AGTR1</i>	rs5186	A	B	B	Moderate
<i>APOA1</i>	rs964184	A	C	C	Weak
<i>APOE</i>	rs439401	A	C	C	Weak
<i>APOB</i>	rs1367117	A	C	C	Weak
<i>CCR5</i>	rs333	B	C	C	Weak
<i>CRP</i>	rs3091244	A	C	C	Weak
<i>DAB2IP</i>	rs7025486	A	A	A	Strong
<i>ELN</i>	rs2071307	B	C	C	Weak
<i>HLA-A</i>	A2	C	C	C	Weak
<i>HLA-B</i>	B61	C	C	C	Weak
<i>HMOX1</i>	GT(n)	C	C	C	Weak
<i>IL6R</i>	rs7529229	A	A	A	Strong
<i>IL10</i>	rs1800896	B	C	B	Weak
<i>LDLR</i>	rs6511720	A	B	A	Moderate
<i>LPA</i>	rs10455872	A	A	A	Strong
<i>LRP1</i>	rs1466535	A	A	A	Strong
<i>MMP2</i>	rs243865	B	C	B	Weak
<i>MMP3</i>	rs3025058	A	A	B	Moderate
<i>MMP9</i>	rs3918242	B	C	C	Weak
<i>MMP13</i>	rs2252070	C	C	C	Weak
<i>MTHFR1</i>	rs1801133	B	C	C	Weak
<i>NOS3</i>	rs1799983	B	C	B	Weak
<i>PLA2G7</i>	rs16874954	B	C	C	Weak
<i>PHACTR1</i>	rs12526453	A	C	C	Weak
<i>SERPINE1</i>	rs1799889	A	C	C	Weak
<i>SORT1</i>	rs599839	A	A	A	Strong
<i>TGFBR1</i>	rs10819634 rs1571590 rs1626340	A	C	C	Weak
<i>TGFBR2</i>	rs1036095 rs764522	A	C	B	Weak
<i>TIMP1</i>	rs4898	C	C	C	Weak
<i>TIMP2</i>	nt573 G/A	C	C	C	Weak
<i>TRIB1</i>	rs2954029	A	C	C	Weak

Figures

Legend to Figure 1: PRISMA flowchart for Systematic Review

Legend to Figure 2: Study frequency and median study size, 1994-2014

Supplementary Tables

Supplementary Table 1: Excluded studies and reasons for exclusion

Supplementary Table 2: Characteristics of included studies

Supplementary Table 3: Number of reports in which genes were investigated in candidate gene studies (including genetic risk score studies)

Supplementary Table 4. Gene regions for which meta-analysis was not possible

Supplementary Table 5. Meta-analyses of odds ratios for associations between genetic variants and AAA.

